

TURFGRASS TOLERANCE TO PREEMERGENCE HERBICIDES

	Bahiagrass	Bermudagrass	Centipede	Kentucky Bluegrass	St. Augustinegrass	Seashore Paspalum	Tall Fescue	Zoysiagrass
PREEMERGENCE								
atrazine	NR	D	T	NR	T	NR	NR	I-T
benefin	T	T	T	T	T	NR	T	T
benefin + oryzalin	T	T	T	NR	T	NR-I	T	T
benefin + trifluralin	T	T	T	T	T	NR-T	T	T
bensulide	T	T	T	T	T	NR-T	T	T
bensulide + oxadiazon	NR	T	NR	T	NR	NR	T	T
dithiopyr	T	T	T	T	T	T	T	T
fenarimol	-	T	-	-	-	-	-	-
isoxaben	T	T	T	T	T	NR	T	T
mesotrione	NR	NR	T	T	I	NR	T	NR
metolachlor	T	T	T	NR	T	NR	NR	T
oryzalin	T	T	T	NR	T	NR-I	T	T
oxadiazon	T	T	NR	T	T	T	T	T
oxadiazon + benefin	NR	T	T	T	T	NR	T	T
pendimethalin	T	T	T	T	T	NR-T	T	T
prodiamine	T	T	T	T	T	T	T	T
pronamide	NR	T	T	NR	T	NR-T	NR	T
simazine	NR	I-T	T	NR	T	NR	NR	T
sulfentrazone + prodiamine	T	T	T	T	NR	T	T	T

T = tolerant; I=intermediate tolerance, use low rates; D = dormant use; NR = not registered for use; S = sensitive

Supplemental 2(ee) label recommendations may be available for some listed pesticides. It is the end-user's responsibility to consult the manufacturer or websites like www.CDMS.net for additional recommendations